## Protein kinase CK2 opens the gate for zinc signaling

Kathryn M. Taylor,<sup>1,\*</sup> Peter Kille<sup>2</sup> and Christer Hogstrand<sup>3</sup>

<sup>1</sup>Breast Cancer Molecular Pharmacology Group; School of Pharmacy and Pharmaceutical Sciences; Cardiff University; Cardiff, UK; <sup>2</sup>Department of Biosciences; Cardiff University; Cardiff, UK; <sup>3</sup>Metal Metabolism Group; Diabetes and Nutritional Sciences Division; King's College London; London, UK

The importance of zinc in biology is evidenced by the growing number of disease states that directly involve aberrant levels of zinc, including neurodegeneration, inflammation, diabetes, cancer and more. Zinc is a component of an estimated 10% of all proteins<sup>1</sup> and has a well-established role in regulation of gene expression, primarily through activation of metalresponsive transcription factor-1 (MTF1) as well as an array of trans-acting factors.2 Recently, attention has been drawn to the fact that free Zn2+ in cells has also non-genomic signaling effects that can be evoked through extracellular stimuli.3,4 The Zn<sup>2+</sup> ion therefore fulfils the criteria for being a second messenger.5 However, delivery of its second messenger role would require three key components to be present: (1) transmission of the extracellular stimuli to the intracellular machinery, (2) mechanism for the gated release of Zn2+ into the cytosol through modification of an effector molecule and (3) the translation of Zn2+ flux into altered activity in downstream targets. Our recent finding that a zinc channel, ZIP7 (SLC39A7), can be activated by phosphorylation by protein kinase CK2 as a direct result of an extracellular stimulation provides the central component of this pathway.6

ZIP7 is the only ZIP zinc channel known to reside in the membrane of the endoplasmic reticulum.<sup>7</sup> In response to epidermal growth factor (EGF) stimulation, CK2 activates ZIP7 through serine phosphorylation, resulting in release of free Zn<sup>2+</sup> ions into the cytosol.<sup>6</sup> The consequential rise in cytosolic Zn<sup>2+</sup> concentration activates protein kinases involved


in cell proliferation, which results in increased cell migration. ZIP7 is ubiquitously expressed in cells unlike many other ZIP channels which are often expressed in a tissue-specific manner. This suggests that ZIP7 may be a gatekeeper for cytosolic Zn2+ release and control Zn2+ signals in multiple cell types.8 The targets directly affected by zinc signals remain to be mapped, but protein tyrosine phosphatases (PTP) seem to be a family of proteins that are particularly affected by an increase in [Zn2+],.4 In a recent study, it was shown that a particular PTP, PTPβ, is inhibited by Zn2+ with a K, of 21 pM, which is well within the range of intracellular Zn2+ concentrations.9 Inhibition of PTPs is thought to be the explanation for the widespread activation of tyrosine kinases observed in cells as a result of elevated levels of Zn2+.5,8 This latter event has major implications for diseases such as cancer that rely on tyrosine kinasedriven signaling pathways for growth and delivers a new and novel area for drug development.

There are still unanswered questions concerning the full extent of the role of zinc in cell signaling. The fact that ZIP7 needs phosphorylation in order to open the gate and release Zn<sup>2+</sup> into the cytosol raises the question of whether other ZIP channels might be gated. There are a number of predicted phosphorylation sites in other zinc transporters and channels<sup>8</sup> that should be investigated. Few of these sites are predicted to be CK2 sites, but several other kinases are implicated, suggesting that different zinc channels may be controlled in different ways, helping

to explain the diversity of zinc signaling functions and the presence of 14 ZIP proteins in humans. There is now a need to identify the direct targets of zinc signals, including how Zn2+ released in the cell can selectively target particular proteins. Protein phosphatases<sup>4,9</sup> and the plasma membrane Ca2+-ATPase10 have been shown to be regulated by Zn2+ at physiologically relevant concentrations, but the exact nature of these interactions remains obscure. Finally, it is also unknown which phosphatase acts to dephosphorylate and presumably inactivate ZIP7, but this discovery may also provide useful avenues for clinical exploration (Fig. 1, see page 1864).

## References

- Passerini A, et al. BMC Bioinformatics 2007; 8:39; PMID:17280606; http://dx.doi.org/10.1186/1471-2105-8-39.
- Hogstrand C, et al. Biochem Soc Trans 2008; 36:1252-7; PMID:19021535; http://dx.doi.org/10.1042/ BST0361252.
- Yamasaki S, et al. J Cell Biol 2007; 177:637-45; PMID:17502426; http://dx.doi.org/10.1083/ jcb.200702081.
- Haase H, et al. Exp Cell Res 2003; 291:289-98; PMID:14644152; http://dx.doi.org/10.1016/S0014-4827(03)00406-3.
- Beavo JA, et al. Nat Rev Mol Cell Biol 2002; 3:710-8; PMID:12209131; http://dx.doi.org/10.1038/nrm911.
- Taylor KM, et al. Sci Signal 2012; 5:11; PMID:22317921; http://dx.doi.org/10.1126/scisignal.2002585.
- Taylor KM, et al. Endocrinology 2008; 149:4912-20; PMID:18583420; http://dx.doi.org/10.1210/en.2008-0351
- Hogstrand C, et al. Trends Mol Med 2009; 15:101-11; PMID:19246244; http://dx.doi.org/10.1016/j.mol-med.2009.01.004.
- Wilson M, et al. J Biol Chem 2012; 287:9322-6; PMID:22275360; http://dx.doi.org/10.1074/jbc. C111 320796
- Hogstrand C, et al. Toxicology 1999; 133:139-45;
  PMID:10378480; http://dx.doi.org/10.1016/S0300-483X(99)00020-7.



**Figure 1.** Schematic illustration of the second messenger signaling pathway proposed for  $Zn^{2+}$ . Various external stimuli cause protein kinase CK2 to phosphorylate endoplasmic reticulum (ER)-located zinc transporter ZIP7, which causes the gated release of  $Zn^{2+}$  stored within the ER through the ZIP7 channel to create a cytosolic "zinc wave." After phosphorylating its target, CK2 dissociates from ZIP7 and may move into the nucleus. We postulate that elevated  $Zn^{2+}$  within the cytosol may selectively inhibit tyrosine phosphatases, leading to prolonged tyrosine kinase activation and an observed downstream increase in pERK and pAKT, resulting in cell proliferation and migration. These events, representing a short-term second messenger activity for Zn, completing within a 20 min time period, are thus temporally separated from established transcriptional impact of zinc ions, which is mediated by metal transcription factor-1 (MTF-1) and is known to regulate metallothionein (MT), glutamylcysteine synthetase (GCS) and ZnT1 (SLC30A1).